

CROP BORON

Fertilizer Group 2
Reg. No. B4693, Act No 36 of 1947

Registration holder: In Line Trading 112 (Pty) Ltd
(Co. Reg. No. 2003/008663/07)
Suite #114, Private Bag X27, Kempton Park, 1620, RSA
Tel: +27 (11) 392 4072
Fax: +27 (11) 392 4363

Active Ingredient (w/w):
Boron (B) 100 g/kg
Wetting Agent 250 g/kg
{Amino-Gluconic Acid™ Chelated}

Crop Boron is a liquid Amino-Gluconic Acid™ Chelated plant nutrient blend, specifically designed to prevent and correct Boron deficiencies which may impair crop growth and reduce yields.

DIRECTIONS FOR USE: Use only as directed.

Compatibility:

Crop Boron is not compatible with herbicides. Never mix **Crop Boron** with alkaline materials including Bordeaux -mixtures and lime sulphur. It is advisable to check the manufacturer recommendations or to do a small miscibility test prior to mixing of **Crop Boron** with other products. **Crop Boron** has a built in wetting agent, thus no additional wetting is required when used.

Mixing:

Shake well prior to opening. Always mix **Crop Boron** with ample quantities of water. Never apply **Crop Boron** undiluted. Follow standard tank mixing procedures.

Storage:

Ensure **Crop Boron** do not freeze. Store in a cool dry place. Never store in direct sunlight for prolonged periods

Application:

Crop Boron should preferably be applied in early morning or late afternoon. Never apply **Crop Boron** to crops undergoing drought stress.

General:

Consult with a qualified agronomist for crop specific information. **Crop Boron** should preferably be applied based on recommendations derived from soil and/or foliar analysis

NOTE: Crop Boron is an Alkaline Product with pH=7.6

Crop / Gewas	Rate /ha Dosing/ha	Application / Toediening
Pome Fruit Stone Fruit Avocado Deciduous Fruit	1 - 2lt	2 Sprays per season, first during early blossoming, repeated again post harvest.
Citrus Litchi Mangoes Macadamia Pecans	50ml / tree in the rootzone or 200ml / 100lt water on foliage	Apply 2 Sprays starting at late flowering (Just before petal fall).
Soya Beans Field Beans Groundnuts Grains	350ml	2 - 4 Sprays at 21 day intervals, prior to and during flowering.
Potatoes Onions	350ml	2 - 3 Sprays during tuber and bulb initiation at 10 day intervals.
Grapes (Table & Wine)	100ml / 100lt 200ml / 100lt	Prior to flowering Post harvest
Cruciferae Vegetables Sunflowers	750ml	Vegetables: 2 - 4 Sprays at 10 day intervals starting 2 - 4 weeks after transplant Sunflowers: 2 - 3 Sprays up to knee height
Tomatoes Tobacco	750ml	2 - 4 Sprays at 10 day intervals prior to and during flower formation and before topping of Tobacco

SHAKE WELL BEFORE USE / SKUD GOED VOOR GEBRUIK