

CROP CAL

Fertilizer Group 2
Reg. No. B4692, Act No 36 of 1947

Registration holder: In Line Trading 112 (Pty) Ltd
(Co. Reg. No. 2003/008663/07)
Suite #114, Private Bag X27, Kempton Park, 1620, RSA
Tel: +27 (11) 392 4072,
Fax: +27 (11) 392 4363

Active Ingredient (w/w): Calcium (Ca) 130 g/kg, Magnesium 1.6 g/kg {Amino-Gluconic Acid™ Chelated}

Crop Cal is a liquid Amino-Gluconic Acid™ Chelated plant nutrient blend, specifically designed to prevent and correct Calcium deficiencies which may impair crop growth and reduce yields.

DIRECTIONS FOR USE: Use only as directed

Compatibility:

Crop Cal is not compatible with herbicides, Phosphate and Sulphate containing products. Never mix **Crop Cal** with alkaline materials including Bordeaux-mixtures and lime sulphur. It is advisable to test prior to mixing of Crop Cal with other products.

Mixing:

Shake well prior to opening. Always mix Crop Cal with ample quantities of water. Never apply **Crop Cal** undiluted. Follow standard tank mixing procedures.

Storage:

Ensure **Crop Cal** do not freeze. Store in a cool dry place. Never store in direct sunlight for prolonged periods.

Application:

Crop Cal should preferably be applied in early morning or late afternoon. Never apply **Crop Cal** to crops undergoing drought stress.

General:

Consult with a qualified agronomist for crop specific information. **Crop Cal** should preferably be applied based on recommendations derived from soil and/or foliar analysis.

Crop / Gewas	Rate /ha Dosering/ha	Application / Toediening
Pome Fruit Stone Fruit Avocado Deciduous Fruit	10lt (250 - 500ml / 100lt)	6 - 12 Sprays per season, starting during early fruit set. For bitterpit sensitive crops use the maximum application rate. Increase the dosage if vigorous shoot growth appears.
Citrus Litchi Mangoes Macadamia Pecans	4 - 10lt 250 - 500ml / 100lt)	Apply 3 - 6 Sprays starting at late flowering (Just before petal fall) at 10 - 14 day intervals.
Soya Beans Field Beans Groundnuts Grains	2 - 3lt	2 Sprays at 7 - 14 day intervals, during and after flowering.
Potatoes Onions	3 - 5lt	3 - 5 Sprays during tuber and bulb initiation at 10 day intervals.
Grapes (Table & Wine)	3 - 5lt	4 - 6 Sprays starting directly after fruit set
Cruciferae Vegetables Sunflowers	750ml	Vegetables: 2 - 4 Sprays at 10 day intervals starting 2 - 4 weeks after transplant Sunflowers: 2 - 3 Sprays up to knee height
Tomatoes Cucurbits Capsicums	3 - 4lt	3 - 5 Sprays at 10 day intervals during fruit formation

SHAKE WELL BEFORE USE / SKUD GOED VOOR GEBRUIK