

# CROP ZINC

Fertilizer Group 2  
Reg. No. B4694 Act 36 (1947)

Registration holder: In Line Trading 112 (Pty) Ltd  
(Co. Reg. No. 2003/008663/07)  
Suite #114, Private Bag X27, Kempton Park, 1620, RSA  
Tel: +27 (11) 392 4072,  
Fax: +27 (11) 392 4363

**Active Ingredient (w/w):**  
**Zinc (Zn) 150 g/kg**  
{Amino-Gluconic Acid™ Chelated}

**Crop Zinc is a liquid Amino-Gluconic Acid™ Chelated plant nutrient blend, specifically designed to prevent and correct Zinc deficiencies which may impair crop growth and reduce yields.**

**DIRECTIONS FOR USE:** Use only as directed

**Compatibility:**

**Crop Zinc** is not compatible with herbicides. Never mix **Crop Zinc** with alkaline materials including Bordeaux-mixtures and lime sulphur. It is advisable to check the manufacturer specifications or to do a small miscibility test prior to mixing of **Crop Zinc** with other products.

**Mixing:**

Shake well prior to opening. Always mix **Crop Zinc** with ample quantities of water. Never apply **Crop Zinc** undiluted. Follow standard tank mixing procedures.

**Storage:**

Ensure **Crop Zinc** do not freeze. Store in a cool dry place. Never store in direct sunlight for prolonged periods.

**Application:**

**Crop Zinc** should preferably be applied in early morning or late afternoon. Never apply **Crop Zinc** to crops undergoing drought stress.

**General:**

Consult with a qualified agronomist for crop specific information. **Crop Zinc** should preferably be applied based on recommendations derived from soil and/or foliar analysis

Crop / Gewas	Rate /ha Doserings/ha	Application / Toediening
Pome Fruit Stone Fruit Avocado Deciduous Fruit	2 - 4lt (50 - 100ml / 100lt)	2 - 4 Sprays starting on early flowering and new growth, repeated at 7 - 14 day intervals. Last application is done post harvest.
Citrus Litchi Mangoes Macadamia Pecans, Coffee	2 - 3lt 50 - 100ml / 100lt)	2 - 4 Sprays on new Spring growth prior to and during flowering.
Soya Beans Field Beans Groundnuts Grains & Maize	2lt (100 - 200ml / 100lt)	2 Sprays early in the season at 7 - 14 day intervals or when deficiency is visible.
Potatoes Onions	2 - 3lt (200ml / 100lt)	2 Sprays at 7 - 14 day intervals, when deficiency is visible.
Grapes (Table & Wine)	2 - 4lt	2 - 4 Sprays at weekly intervals during active growth period
Vegetables Capsicums Tomatoes	2 - 3lt	2 Sprays at 7 - 14 day intervals, when deficiency is visible.
Cucurbits Crucifers	3lt	2 - 3 Sprays at 10 day intervals when deficiency is visible

**SHAKE WELL BEFORE USE / SKUD GOED VOOR GEBRUIK**