

# NUTRISPRAY N

Fertilizer Group 1  
Reg. No. K8789, Act No. 36 of 1947

Registration holder: In Line Trading 112 (Pty) Ltd  
(Co. Reg. No. 2003/008663/07)  
Suite #114, Private Bag X27, Kempton Park, 1620, RSA  
Tel: +27 (11) 392 4072,  
Fax: +27 (11) 392 4363

Nitrogen / Stikstof (N).....150 g/kg  
Copper / Koper (Cu).....2.6 g/kg  
Boron / Boor (B).....11.5 g/kg  
Molybdenum / Molibdeen (Mo).....4 g/kg

SG = 1.2 @ 20°C

## WARNING

**This product contains Boron. The misuse of NUTRISPRAY N can cause serious damage to crops. Use this product according to the recommendations on the label.**

## GENERAL:

- \* **NUTRISPRAY N** is a water-soluble foliar feed solution, which is readily available with a wide range of nutrients for plants.
- \* Particularly suitable for foliar sprays, as well as for applications through irrigation systems.

## DIRECTIONS FOR USE : Use only as directed.

- \* **NUTRISPRAY N** should be used as a supplement to normal fertilizing practices, especially during periods of nutrient deficiency.
- \* **NUTRISPRAY N** should be applied to established plants as a full cover spray during periods of high nutrient demands, such as prior to flowering, fruit and after fruit set.
- \* The application rate depends on the soil fertility and reaction needed. The lower dosages should be applied for maintenance and the higher dosage rates nutrient deficiencies.
- \* Application may be performed at 7- to 14-day intervals depending on growing conditions.
- \* The recommended amount of **NUTRISPRAY N** should be added to sufficient water for a full cover spray of one hectare, unless otherwise stated and is suitable for use in all types of spray equipment.
- \* **NUTRISPRAY N** should be applied in the early morning or late afternoon, when leaf tissue contains maximum moisture and there are no moisture- and temperature stresses.
- \* Consult the distributor, consultant and/or registration holder if crop specific programmes are followed using this product.

## Compatibility :

**NUTRISPRAY N** is not compatible with alkaline chemicals and products with high calcium content. Do not apply in conjunction with Copper fungicides, especially Copper Oxochlorides, and Lime Sulphurs. It is advisable to check the manufacturer recommendations or to do a small miscibility test prior to mixing **NUTRISPRAY N** with other chemicals.

## Application Rate:

Apples, Pears, Stone Fruit: 200ml - 500ml /100l spray water, at maximum 5l / ha. Apply 2 - 3 times at 10 - 14 day intervals prior to harvest.  
Deciduous Fruit: 200ml / 100l. Apply 2 times at 7 day intervals during bud break.  
Barley, Oats, Wheat: 2l / ha. Apply once at growth stage 7, repeat at stage 13 if required.  
Carrots: 2l / ha. Apply 2 - 3 times at 7 - 10 day intervals from 15 cm crop length.  
Citrus: 2l / ha. Apply 2 - 3 times between bud break and 100% flowering.  
Maize: 2l / ha at 4 - 8 leaf stage.  
Onions: 2l - 4l / ha. Apply 2 times at 10 - 14 day intervals during bulb swelling.  
Potatoes: 2 - 4l / ha. Apply 2 - 3 times at 10 - 14 day intervals a month after germination.  
Vines: 2l / ha, repeated 2 - 3 times at 10 - 14 day intervals from 10 - 15cm shoot length.