

# Amino Gluconate CalMag + B

FERTILISER GROUP 2  
Reg. No.B3704, Act 36 of 1947

Calcium (Ca)..... 73 g/kg  
Magnesium (Mg) ....28 g/kg  
Boron (B) .....3.5 g/kg  
{Amino-Gluconic Acid™ Chelated}  
SG - 1.38

**Amino Gluconate CalMag+B is a liquid Amino-Gluconic Acid Chelated plant nutrient blend, specifically designed to prevent and correct Calcium & Magnesium deficiencies which may impair crop growth and reduce yields.**

**DIRECTIONS FOR USE:** Use only as directed

**Compatibility:**

**Amino Gluconate CalMag+B** is compatible with most agricultural products, including insecticides, fungicides & nutrients. **Amino Gluconate CalMag+B** is not compatible with herbicides, Sulphates and certain Phosphates. Never mix **Amino Gluconate CalMag+B** with alkaline materials including Bordeaux-mixtures and lime sulphur. It is advisable to do a miscibility test prior to mixing of **Amino Gluconate CalMag+B** with other products.

**Mixing:**

Shake well prior to opening. Always mix **Amino Gluconate CalMag+B** with ample quantities of water. Never apply **Amino Gluconate CalMag+B** undiluted. Follow standard tank mixing procedures.

**Storage:**

Ensure **Amino Gluconate CalMag+B** do not freeze. Store in a cool dry place. Never store in direct sunlight for prolonged periods

**Application:**

**Amino Gluconate CalMag+B** should preferably be applied in early morning or late afternoon. Never apply **Amino Gluconate CalMag+B** to crops undergoing drought stress.

**General:**

Consult with a qualified agronomist for crop specific information. **Amino Gluconate CalMag+B** should preferably be applied based on recommendations derived from soil and/or foliar analysis

**Directions for use**

(Use only as directed and shake well before use)

The pH of the spray solution must preferably be between 5.0 and 6.5. pH correction must be done with a pH-corrector suitable with foliar nutrients

Crop/Gewas	Rate /ha Dosering /ha	Application / Toediening
Pome Fruit Stone Fruit Avocado Deciduous Fruit	10lt	6 - 12 Sprays starting directly after fruitset at 10 day intervals. Use higher application on Bitterpit sensitive crops.
Citrus Litchi Mangoes Macadamia Pecans	10lt	4 - 6 Sprays starting during late flowering (Just before petal fall) at 10 day intervals. Add WETTA as wetting agent
Soya Beans, Grains Field Beans, Groundnuts	3lt	2 - 3 Sprays at 10 day intervals. Add WETTA as wetting agent
Potatoes	3lt	3 - 5 Sprays starting at tuber initiation at 10 day intervals.
Grapes (Table & Wine)	4lt	4 - 6 Sprays directly after fruitset at 10 day intervals. Add WETTA
Cruciferae Vegetables Cotton	2lt	2 - 4 Sprays at 10 day intervals starting 2 - 4 weeks after transplant

Registration holder: In Line Trading 112 (Pty) Ltd  
(Co. Reg. No. 2003/008663/07)  
Suite #114, Private Bag X27, Kempton Park, 1620, RSA  
Tel: +27 (11) 392 4072.  
Fax: +27 (11) 392 4363

SKUD GOED VOOR GEBRUIK • SHAKE WELL BEFORE USE