

CALMAG+B

Fertilizer Group 2
Reg. No. B4696, Act No 36 of 1947

Registration holder: In Line Trading 112 (Pty) Ltd
(Co. Reg. No. 2003/008663/07)
Suite #114, Private Bag X27, Kempton Park, 1620, RSA
Tel: +27 (11) 392 4072.
Fax: +27 (11) 392 4363

Active Ingredient (w/w):
Calcium (Ca) 73 g/kg Magnesium (Mg) 28 g/kg
Boron (B) 3.5 g/kg
{Amino-Gluconic Acid™ Chelated}

Crop CalMag + B is a liquid Amino-Gluconic Acid™ Chelated plant nutrient blend, specifically designed to prevent and correct Calcium, Magnesium & Boron deficiencies which may impair crop growth and reduce yields.

DIRECTIONS FOR USE: Use only as directed

Compatibility:

Crop CalMag+B is not compatible with herbicides, Phosphate and Sulphate containing products. Never mix **Crop CalMag+B** with alkaline materials including Bordeaux mixtures and lime sulphur. It is advisable to check the manufacturer specifications or to do a small miscibility test prior to mixing of **Crop CalMag+B** with other products.

Mixing:

Shake well prior to opening. Always mix **Crop CalMag+B** with ample quantities of water. Never apply **Crop CalMag+B** undiluted. Follow standard tank mixing procedures.

Storage:

Ensure **Crop CalMag+B** do not freeze. Store in a cool dry place. Never store in direct sunlight for prolonged periods.

Application:

Crop CalMag+B should preferably be applied in early morning or late afternoon. Never apply **Crop CalMag+B** to crops undergoing drought stress.

General:

Consult with a qualified agronomist for crop specific information. **Crop CalMag+B** should preferably be applied based on recommendations derived from soil and/or foliar analysis.

| Crop / Gewas | Rate /ha Dosering/ha | Application / Toediening |
|---|----------------------------------|--|
| Pome Fruit Stone Fruit Avocado Deciduous Fruit | 10lt (250 - 500ml / 100lt) | 6 - 12 Sprays per season, starting during early fruit set. For bitterpit sensitive crops use the maximum application rate. Increase the dosage if vigorous shoot growth appears. |
| Citrus Litchi Mangoes Macadamia Pecans | 4 - 10lt 250 - 500ml / 100lt) | Apply 3 - 6 Sprays starting at late flowering (Just before petal fall) at 10 - 14 day intervals. |
| Soya Beans Field Beans Groundnuts Grains | 2 - 3lt | 2 Sprays at 7 - 14 day intervals, during and after flowering. |
| Potatoes Onions | 3 - 5lt | 3 - 5 Sprays during tuber and bulb initiation at 10 day intervals. |
| Grapes (Table & Wine) | 3 - 5lt | 4 - 6 Sprays starting directly after fruit set |
| Cruciferae Vegetables Sunflowers | 750ml | Vegetables: 2 - 4 Sprays at 10 day intervals starting 2 - 4 weeks after transplant Sunflowers: 2 - 3 Sprays up to knee height |
| Tomatoes Cucurbits Capsicums | 3 - 4lt | 3 - 5 Sprays at 10 day intervals during fruit formation |

SHAKE WELL BEFORE USE / SKUD GOED VOOR GEBRUIK